

P: 734-429-9053 | F: 734-944-3934

968 E Michigan Ave | Saline, MI 48176

Mon - Fri: 9am-6pm | Sat: 9am-3pm | Sun: Closed

COMPOUNDED SEMAGLUTIDE

PATIENT INFORMATION	PRESCRIBER INFORMATION
Name:	Name:
DOB:	NPI:
Address:	Address:
Phone:	Phone:
Allergies:	Fax:
PRESCRIPTION (SELECT ONE)	
13-Week Protocol : This protocol is based on p maintenance dose efficiently.	patient feedback and response to get patients to an effective
1st fill: Semaglutide 1mg/ml Suspension Place 0.5ml under the tongue for a minimum of 2 minute eat or drink anything for 30 minutes after dose	es once daily for 2 weeks then increase to 1ml once daily. Do not
2nd fill: Semaglutide 2mg/ml Suspension Place 1ml under the tongue for a minimum of 2 minutes	s once daily. Do not eat or drink anything for 30 minutes after dose
3rd fill: Semaglutide 3mg/ml Suspension Place 1ml under the tongue for a minimum of 2 minutes	s once daily. Do not eat or drink anything for 30 minutes after dose
Maintain dosage of 3mg/ml daily thereafter	
Other: Once daily, placeml under tong drink for 30 minutes after dose.	gue for a minimum of 2 minutes before swallowing. Do not eat or
SIG: Compounded Semaglutide in SubMagna ^{TI}	M HMW (Select one)
1mg/ml suspension	2mg/ml suspension3mg/ml suspension
Oty (in 30ml increments):	
Refill times	
Prescriber's Signature:	Date: